



POST-OPERATIVE ORDERS ARTHROSCOPIC CORACOCLAVICULAR RECONSTRUCTION

DAY OF SURGERY:

Because of your Anesthesia

- Do Not operate hazardous machinery for 24 hours.
- Do Not make important personal or business decisions or sign legal documents for 24 hours.
- Have someone stay with you for the first 24 hours post-operatively.
- You will be sleepy most of the day. Get some rest.
- Advance your diet as tolerated.
- Avoid alcohol for the first 24 hours.
- Your interscalene block will begin to wear off after several hours. Be careful to support your arm if removing from the sling, as you may not have independent control.

Medication Information

- A prescription for a pain medication may be prescribed for you. This medication is a controlled substance and is meant to be taken AS NEEDED for pain only. These medications can be addictive if taken over long periods of time and should be used thoughtfully to control your pain given their known risk of dependence and addiction. If you cannot tolerate your medication and feel you need something greater than an over the counter medication, call the office. DO NOT take this medication in combination with Tylenol.
 - Tramadol - 1-2 pills every 6 hours AS NEEDED for pain
 - Tylenol #3 - 1-2 pills every 6 hours AS NEEDED for pain
 - Norco 5/325 1-2 pills every 4-6 hours AS NEEDED for pain
 - Percocet 5/325 1-2 pills every 4-6 hours AS NEEDED for pain
 - Nucynta 50 mg 1-2 pills every 6 hours AS NEEDED for pain
- Avoid taking anti-inflammatory medication for the next 3 weeks as these medications can interfere with healing. Anti-inflammatory medications include Ibuprofen, Naproxen, Aleve, Meloxicam, and Celebrex.
- If taking your pain medication regularly, you should also take two protective stomach medications during this time period. You may choose these medications from the list below.
 - Colace 100 mg twice a day
 - Protonix 40 mg daily
 - Nexium 20 mg daily
 - Prilosec OTC 20 mg daily
- Because of concerns due to inactivity following surgery, please adhere to the following instructions;
 - You should take an aspirin daily 325 mg
 - Please wear your compression stockings until you are fully mobile
 - Ambulate frequently and pump your ankles/feet often to encourage blood flow

Activity

- Keep your arm in the sling at all times.
- Elevate arm when sitting and resting. This will prevent swelling and decrease pain
- Ambulate as tolerated
- Sleeping in a recliner may also be more comfortable for the first night or two. Additionally, if lying down, elevate your head and place a small pillow under your arm to prevent it from falling back.

Dressing

- Keep dressing intact, clean and dry.
- Cold therapy is used to assist in decreasing pain and swelling. Whatever cold/ice method is used, keep it on for 20-30 minutes at a time with breaks of 60-90 minutes. Always avoid direct application to the skin if using ice in a plastic bag.



DAY AFTER SURGERY:

- If you have not done so, please call our office to schedule your follow up appointment 7-10 days following your surgery yesterday. The office number is (636) 778-3177.
- You may remove your dressing today, however be careful to leave the white “steri-strips” on the skin. Do Not apply any creams, ointments, balms or other material to the incision or around it. Do Not swim or take baths until wounds are completely healed.
 - You may shower today. Do not scrub the incisions, but you may allow the water to run over them. Otherwise, keep the incisions covered. To wash under the arm, bend over at the waist and let the arm passively move over away from the body.
 - REMAIN in the sling at all times except for hygiene. Keep the arm supported when performing hygiene so the weight of the arm does not pull on the repair/reconstruction

TWO DAYS POST OP...UNTIL POST OP OFFICE VISIT:

Continue all other instruction as listed previously.

- A healthy and nutritious diet including low fat and high protein is encouraged to promote healing of tissues
- Frequent ambulation daily is encouraged to maintain strength and endurance in your legs and trunk.
- Avoid trying to use the narcotic medications for sleep as narcotic induced sleep is not restorative and pre-disposes to dependence
- Avoid ANY tobacco or nicotine use – This significantly reduces the likelihood of a good outcome and pre-disposes you to INFECTION and decreases blood flow to the healing area.

Please call if you develop a nausea, vomiting or a rash after taking your medication.

Call if you develop a fever of greater than 100 degrees, chills, any redness, swelling, bleeding or pus-like drainage coming from the wound, or if the wound opens up.

Call if your pain seems excessive or if you have any questions or concerns. Our office number is (636) 778-3177.